

Charmonium and lepton pair production with medium energy antiprotons *

Boris Kopeliovich

*Max-Planck Institut für Kernphysik, Postfach 103980, 69029 Heidelberg, Germany
& Joint Institute for Nuclear Research, Dubna, 141980 Moscow Region, Russia*

Abstract

The medium energy antiproton beam facility which is under discussion at GSI offers new opportunities to study the QCD of heavy flavours, nuclear structure and exotic mesons. We discuss these topics and briefly highlight other problems in charm production and proton structure.

1 Charmonium production off nuclei

1.1 Properties of charmonia

Charmonium spectroscopy is known as a laboratory for heavy quark QCD. Realistic potential models [1] successfully describe the mass spectrum. Not much is known, however, about charmonium properties. Even the cross section of interaction with a nucleon, $\sigma_{tot}(\Psi N)$, is not measured. One desperately needs to know it in order to interpret the data on charmonium production in relativistic heavy ion collisions.

The sources of information about $\sigma_{tot}(J/\Psi N)$ are the photoproduction data on a proton and hadroproduction off nuclei [2]. The former, analysed with VDM gives $\sigma_{tot}(J/\Psi N) = 1.2 mb$ at the SPS energy $\sqrt{s_\Psi} = 10 GeV$, while the latter analysed within the optical model results in $\sigma_{tot}(J/\Psi N) \approx 6 mb$. Data for $\Psi'(2S)$ production in both cases also lead to a puzzling conclusion that $\sigma_{tot}(\Psi' N) \approx \sigma_{tot}(J/\Psi N)$ contradicting the expectation that Ψ' having about twice a big radius as J/Ψ [1] has to interact much stronger. These controversies probably mean that the one-channel approach fails and the higher charmonium excitations

*The talk given at the meeting "Perspectives of Hadron Physics", GSI, Darmshtadt, June 8, 1998

must come to play [2]. The key parameters which control the solution of the two-channel problem for charmonium production off nuclei are

$$\epsilon = \frac{\langle \Psi' | \hat{\sigma} | J/\Psi \rangle}{\langle J/\Psi | \hat{\sigma} | J/\Psi \rangle} \quad (1)$$

$$r = \frac{\langle \Psi' | \hat{\sigma} | \Psi' \rangle}{\langle J/\Psi | \hat{\sigma} | J/\Psi \rangle} \quad (2)$$

$$R = \frac{\sigma(hN \rightarrow \Psi' X)}{\sigma(hN \rightarrow J/\Psi X)} \quad (3)$$

One can only roughly estimate these parameters from data on charmonium hadroproduction off nuclei since the production dynamics is poorly known. A better information could be extracted from virtual photoproduction, but no data are available and are not expected in the near future.

Direct production $\bar{p}p \rightarrow J/\Psi(\Psi')$ has an advantage of known dynamics. besides, it provides the lowest energy of produced charmonium, $p_L(J/\Psi) = 4.06 \text{ GeV}$, $p_L(\Psi') = 6.2 \text{ GeV}^\dagger$. At this energy the contribution from other channels is tiny and one can reliably measure the Ψ -nucleon absorption cross section.

The production cross section off nuclei at the energy of the peak is, however, strongly suppressed compared to $\bar{p}p \rightarrow \Psi$ due to Fermi motion by a factor [3],

$$\frac{\sigma(\bar{p}A \rightarrow \Psi A')}{Z \sigma(\bar{p}p \rightarrow \Psi)} = \pi \frac{M_N}{M_\Psi} \frac{\Gamma_\Psi}{k_F}, \quad (4)$$

where k_F is the mean value of the Fermi momentum. If the beam momentum resolution Δp_b exceeds Γ_Ψ (most probably), then Γ_Ψ should be replaced by Δp_b [3] what may substantially enhance the ratio in (4). This does not mean, however, that with a worse resolution Δp_b the nuclear cross section is larger. The ratio (4) grows just due to decreasing denominator $\sigma(\bar{p}p \rightarrow \Psi)$. At $\Delta p_b = 1 \text{ MeV}$ the production rate off nuclei is suppressed by three orders of magnitude.

The dependence of J/Ψ and Ψ' production cross section on the beam momentum can be evaluated using the Gaussian form for the Fermi momentum distribution with $k_F = 200 \text{ MeV}$. Fig. 1 shows such a dependence in arbitrary units, however, the relative Ψ' to J/Ψ production rate is fixed by

$$\frac{N_{\Psi'}}{N_{J/\Psi}} = \frac{\Gamma_{\bar{p}p}^{\Psi'}}{\Gamma_{\bar{p}p}^{J/\Psi}} e^{-\Delta\sigma \langle T \rangle}, \quad (5)$$

where $\langle T \rangle$ is the mean value of the nuclear thickness function, $\Delta\sigma = \sigma_{tot}(\Psi' N) - \sigma_{tot}(J/\Psi N) \approx 9 \text{ mb}$.

Fig. 1 shows that in the beam momentum range $p_L = 5 - 6 \text{ GeV}$ one can produce either J/Ψ or Ψ' at the same beam momentum. As a result, these two states strongly interfere, leading to a very much modified transparency for the charmonia. The value of the parameter R varies with p_L from zero to infinity, and either J/Ψ or Ψ' can be enhanced by the nucleus,

[†]this is much lower energy than can be achieved in practice in experiments with inverse kinematics at SPS. Even at $x_F = -0.5$ the J/Ψ momentum is 9 GeV .




Figure 1: *Production rate of J/Ψ and Ψ' in $\bar{p} Pb$ annihilation as a function of the beam momentum.*

rather than suppressed. Indeed, the effective "absorption" cross section may be negative dependent on R [4], since $\epsilon < 0$,

$$\begin{aligned}\sigma_{abs}^{J/\Psi} &= \sigma_{tot}(J/\Psi N) \left[1 + \epsilon R F_A^2(q_f) \right] < 0 \quad \text{if } R \gg 1 \\ \sigma_{abs}^{\Psi'} &= \sigma_{tot}(\Psi' N) \left[1 + \frac{\epsilon}{r R} F_A^2(q_f) \right] < 0 \quad \text{if } R \ll 1 .\end{aligned}\quad (6)$$

Here the nuclear form factor depends on the longitudinal momentum transfer related to the formation length [4],

$$q_f = \frac{M_{\Psi'}^2 - M_{J/\Psi}^2}{2 p_L} . \quad (7)$$

Note that even if data on virtual photoproduction of charmonia were available, they would have quite a restricted range of variation of the parameter $0.5 < R < 1$ corresponding to $Q^2 = 0$ and $Q^2 \rightarrow \infty$, respectively.

1.2 J/Ψ probe for nuclear structure

When the beam momentum is far from the overlap region in Fig. 1 the interference effects vanish and the kinematics is certain. This makes J/Ψ production a perfect tool to study Fermi distribution of bound protons, especially its high-momentum tail. Recently experiment E850 at BNL [5] on $A(p, 2p)A'$ with detection of the recoil neutron found a strong correlation between the disbalance of the transverse momenta of the protons and the neutron momentum. This is the first solid evidence in favour of hypothesis that the high-momentum tail of Fermi distribution is due to two-nucleon correlations, rather than to the mean nuclear field.

Such a reaction, however, is unable to measure the longitudinal component of the Fermi momentum because of color transparency effect [6]. On the other hand, deep-inelastic lepton scattering is sensitive only to the longitudinal component. J/Ψ production in $\bar{p}$ annihilation allows to measure both k_L and k_T with a high precision, provided that the beam momentum is known with a high resolution.

The correlation between the longitudinal and transverse components of the Fermi momentum dictated by the light-cone nuclear dynamics is quite different from the intuitive expectation: $\langle k_T^2 \rangle$ grows with k_L . Indeed, a bound nucleon carries a fraction of the total nuclear light-cone momentum, $\alpha = 1/A + \Delta_\alpha$, where Δ_α is related to longitudinal Fermi motion. The energy denominator has a form,

$$\frac{k_T^2}{\alpha(1-\alpha)} + \frac{M_N^2}{\alpha} + (A-1)^2 \frac{M_N^2}{\alpha} - A^2 M_N^2 = \frac{k_T^2 + A^2 M_N^2 \Delta_\alpha^2}{\alpha(1-\alpha)}. \quad (8)$$

Therefore,

$$\langle k_T^2 \rangle = A^2 M_N^2 \Delta_\alpha^2 = k_L^2. \quad (9)$$

Thus, we arrived at quite an unusual correlation: the mean value of k_T^2 grows with k_L . Note that this correlation is independent of the origin of high Fermi momenta, mean field or two-nucleon correlation. To disentangle between them in J/Ψ production by antiprotons one should detect a recoil neutron, like is done in [5].

2 Search for the 4-quark mesons in Drell-Yan reaction

A diquark, *i.e.* a qq pair in an antitriplet color state is different from $\bar{q}$ only by spin and mass, unless the interaction is hard enough to resolve its qq structure. Replacing the $\bar{q}$ in a meson by a diquark one gets a baryon. It looks very natural to replace the quark in a baryon by antidiquark. Then one arrives at a 4-quark meson having a diquark-antidiquark structure, $D - \bar{D}$. One of the candidates for such a state is the $C(1480)$ meson, found in the OZI suppressed decay mode $\phi\pi^0$ [7] (see also in [8]).

A $\bar{p}p$ collision is a natural source for such 4-quark mesons, since the diquark and antidiquark already pre-exist. One should annihilate the beam-target valence $\bar{q} - q$ pair and the rest is an excited $\bar{D}D$ pair, as is shown in Fig. 2a. The effective mass of $\bar{D}D$ is, however, larger than $2M_N$ and a decay to a lighter 4-q state is difficult. Indeed, decay of the color string as it is shown in Fig. 2a leads to a production of a baryon-antibaryon pair and mesons.

An effective way to form a desired 4-quark state is to produce a lepton pair via Drell-Yan mechanism as is shown in Fig. 2b. First of all, the lepton pair signals that the $\bar{q}q$ annihilation did occur. Secondly, it takes away an extra energy and the rest 4-quark state may have any mass. Thirdly, the lepton pair is easy to detect and it fixes the missing mass of the 4-quark system, provided that the lepton momenta are well measured. One can also detect the decay products of the 4-quark meson which would help much in establishing its quantum numbers.

This reaction suggested in [9] can be studied complementary to charmonium production. The preferred effective mass interval is $1.5 < M_{e^+e^-} < 3 \text{ GeV}$. The reaction was simulated in [9] for kinematics corresponding to the geometry of the SuperLEAR internal beam detector,




Figure 2: a) $\bar{p}p$ interaction with annihilation of a pair of projectile valence $\bar{q}q$ pair. The produced diquark-antidiquark state mass is $M_{\bar{D}D} > 2M_N$. b) The same as in fig. 2a, but accompanied with production of a Drell-Yan lepton pair. A 4-quark state of any mass can be produced.

and the cross section parameterized as,

$$\frac{d\sigma(\bar{p}p \rightarrow e^+e^- X)}{dM_{e^+e^-}} = \frac{0.21}{M_{e^+e^-}^3} e^{-11.7\tau} (\mu b) , \quad (10)$$

where $\tau = M_{e^+e^-}^2/s$ and $M_{e^+e^-}$ is in GeV .

It was found in [9] that with luminosity $10^{32} cm^{-2}sec^{-1}$, the overall efficiency 0.5 and the trigger efficiency 0.7 one can expect ~ 1200 events a day with a $12 GeV$ antiproton beam. For a light 4-quark system, $0.9 < M_{4q} < 2 GeV$, the expected rate is ~ 300 events a day. For this mass interval the production rate is maximal for the beam momentum range $5 - 8 GeV$.

3 What else?

3.1 Open charm production

- *Inclusive production of leading D and $\bar{D}$ mesons.* $D(\bar{D})$ mesons (not to be mixed up with diquarks) are produced in $\bar{p}p \rightarrow D(\bar{D})X$ at high x_F in a very unusual configuration: the major fraction of the longitudinal momentum is carried by the projectile light quark. In this configuration the D meson is effectively stopped by interaction with nuclear matter, because after the D meson breaks up the slow c quark produces a D meson with low x_F . This is very different from propagation of a formed D meson containing a leading c quark, which loses only a tiny fraction of its momentum interacting with medium. Nuclear suppression of D meson production rate versus Feynman x_F would provide precious information about space-time pattern of formation of the D meson wave function.

- *Exclusive production $\bar{p}p \rightarrow D\bar{D}$.* In this case a very strong leading effect is expected: the $D(\bar{D})$ mesons are predominantly produced in the fragmentation regions of $\bar{p}(p)$. When this reaction occurs in a nuclear environment, nuclear suppression of production rate serves as a direct measurement of D and $\bar{D}$ absorption via inelastic interactions with bound nucleons on the way out of the nucleus. As different from inclusive D meson production, the DN and $\bar{D}N$ inelastic cross sections can be measured.

3.2 Proton structure and nature of baryon number

- *Quark intrinsic momentum in a diquark.* Comparative measurement of transverse momenta of pions produced in annihilation and non-annihilation channels of $\bar{p}p$ interaction provides unambiguous information about the diquark size. In the former case the (anti)diquark in the (anti)proton breaks up and releases the intrinsic quark momenta, which are expected to be much larger than what we usually observe in inelastic hadronic collisions. Indeed, there are many indications [10], both experimental and theoretical, that the diquark in a proton has a size much smaller than the proton radius. We suggest to use this as a very sensitive way to measure the quark Fermi-momentum inside the diquark. A strong $u - d$ quark correlation is expected due to the gluonic structure of the QCD vacuum (instanton model [11]) and even in potential models [12].
- *Mechanisms of baryon stopping* is a hot topic in physics of relativistic heavy ion collisions [13]. One can study this in more detail in $\bar{p}p$ annihilation where the baryon number flows over the whole rapidity interval. According to the modern classification [14, 15] baryon number can be transferred by the diquark (important only at rather low energy, up to a few GeV [16]), by a valence quark plus string junction [17] (mid energies) and by a pure gluon field [16, 18, 19, 20] (=string junction). These mechanisms have a very different multiplicity of produced particles, 1:2:3, respectively. One can effectively disentangle these three mechanisms by analyzing the multiplicity distribution in annihilation events [19]. One can also test the so-called Eylon hypothesis [21, 15] that annihilation is related through the unitarity relation to the Pomeron, i.e. it does not contribute to the $\bar{p}p$ and pp total cross section difference.
- Antiproton production with high p_T in $\bar{p}p$ interaction is known to be due to high- p_T scattering of the projectile antidiquark. Therefore, a high- p_T antiproton produced in $\bar{p}A$ collision should attenuate due to annihilation of the diquark propagating through nuclear matter [22]. The annihilation (break up) cross section strongly depends on the diquark size (an analogue to color transparency). Therefore one can get precious information about the diquark radius [22].

Summarizing, a high luminosity antiproton beam at GSI would open new opportunities for study of QCD dynamics of strong interactions and hadronic structure. Many of them are unique. This talk covers only a small part of the possibilities particularly presented by other speakers at this meeting.

Acknowledgements: I am grateful to Jörg Hüfner and Wolfram Weise for collaboration in formulation of the problems presented in the first two sections. The ideas proposed for

experimental study in the last section were generated during stimulating discussion with A. Gillitzer, S. Paul and W. Weise. I am also thankful to Sverker Fredriksson who read the paper and made valuable comments and Josef Pochodzalla for useful discussions.

References

- [1] W. Buchmüller and S.-H. H. Tye, Phys. Rev. **D24** (1981) 132
- [2] J. Huefner and B.Z. Kopeliovich, Phys. Lett. **B426** (1998) 154
- [3] G.R. Farrar, L.L. Frankfurt, M.I. Strikman, and H. Liu, Nucl. Phys. **B35** (1990) 125
- [4] J. Huefner and B.Z. Kopeliovich, Phys. Rev. Lett. **76** (1996) 192
- [5] The E850 Collaboration, J. Aclander et al., nucl-ex/9801003
- [6] B.K. Jennings and B.Z. Kopeliovich, Phys. Rev. Lett. **70** (1993) 3384
- [7] S.I. Bityukov et al., Phys. Lett. **B188** (1988) 383
- [8] B.Z. Kopeliovich and E. Predazzi, Phys. Rev. **D51** (1995) 2114
- [9] A.V. Bannikov, Yu.A. Gornushkin, B.Z. Kopeliovich, Z.V. Krumshtein and M.G. Sapozhnikov, preprint JINR, E1-92-344, Dubna, 1992 (is available in the KEK library: http://ccdb1.kek.jp/cgi-bin/img_index?9302455)
- [10] M. Anselmino et al., Rev. Mod. Phys. **65** (1993) 1199
- [11] M.M. Musakhanov, P.K. Choudhuri, V.P. Isaev, Yad. Fiz. **45** (1987) 790
- [12] S. Fredriksson, J. Hansson and S. Ekelin, Z. Phys. **C75** (1997) 107
- [13] A. Capella and B.Z. Kopeliovich, Phys. Lett. **B381** (1996) 325
- [14] G.C. Rossi and G. Veneziano, Nucl. Phys. **B123** (1977) 507; Phys. Rep. **63** (1980) 149
- [15] B.Z. Kopeliovich and B.G. Zakharov, Sov. Phys. Particles and Nuclei, **22** (1991) 140
- [16] B.Z. Kopeliovich, Sov. J. Nucl. Phys. **45** (1987) 1078
- [17] B.Z. Kopeliovich and B.G. Zakharov, Sov. J. Nucl. Phys. **49** (1989) 674
- [18] B.Z. Kopeliovich and B.G. Zakharov, Sov. J. Nucl. Phys. **48** (1988) 136
- [19] B.Z. Kopeliovich and B.G. Zakharov, Phys. Lett. **211B** (1988) 221
- [20] B.Z. Kopeliovich and B.G. Zakharov, Z. Phys. **C43** (1989) 241
- [21] Y. Eylon and H. Harari, Nucl. Phys. **B80** (1974) 349
- [22] B.Z. Kopeliovich and B.G. Zakharov, Czech. J. Phys. **41** (1991) 409